



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

of which I give in the following extract from Saunier's "Traité D' Horlogerie Moderne."

"M. Phillips a démontre', dans un Mémoire présenté a l' Académie des sciences, que l' étendue des oscillations du balancier, ou' le défaut d' équilibre s'apercevra le moins au réglage dans les différentes positions inclinées, est de $439^{\circ}28'$, soit en nombre rond 440° , un peu moins de 1 tour $\frac{1}{4}$."

Whether this demonstration was analytical or experimental I cannot say, not having seen the Memoir referred to. In the course of twenty years experience in rating and adjusting watches and pocket chronometers, many opportunities have occurred to verify Prof. Phillips' conclusion.

The arc mentioned seems to be independent of the amount of error of poise, as experiments with a balance only slightly out of poise, and with the same balance after removing a whole screw from one portion of its rim will show.

In a marine chronometer, the balance vibrates in a horizontal plane, and is not so susceptible to errors of poise, but if the balance is made to vibrate as near to this arc as possible, any induced polarity, caused by the magnetism of iron ships, will not cause so much irregularity in the rate as it would if the balance had either a larger or smaller arc of vibration.

J. M. ARNOLD.

Boston, Mass.

FURTHER REMARKS ON THE SOLUTION OF PROBLEM 260, BY THE EDITOR.—The solution of problem 260, given at pp. 121–122, as corrected by the Note at p. 151, is, we trust, a sufficient discussion of that question, as announced at p. 95. It will be seen, however, that equations (2) and (3) as written at the top of p. 122 (with a slight alteration of signs), are adapted to the solution of a different question, viz., Prob. 8, ANALYST, Vol. I, page 35.

Writing $R + r$ instead of $R - r$ in (2) and changing all the signs in (3), equation (1) retaining the negative sign, as written at p. 121, we have, for the equivalent of (6) on p. 122,

$$\left(\frac{d\varphi}{dt}\right)^2 = \frac{10}{7} \frac{g}{R+r} (1 - \cos \varphi); \quad (6)$$

and writing $R' = 0$ in (3) it becomes

$$\left(\frac{d\varphi}{dt}\right)^2 = \frac{g}{R+r} \cos \varphi. \quad (7)$$

Equating the right hand members of (6) and (7) and reducing, we get $\cos \varphi = \frac{1}{4}$. (See foot-note, p. 35, ANAL. Vol. I.)

If, instead of quitting the sphere when $R' = 0$, the rolling sphere is constrained to move on the surface of the fixed sphere until $\cos \varphi = 0$, then will the solution of (6) give the time t , in which the rolling sphere will descend from the summit of the fixed sphere over one fourth of its circumf., viz.;

$$t = \frac{1}{\sqrt{a}} \left[\frac{2}{\sqrt{2}} \log \frac{\sqrt{1+\cos \varphi} - \sqrt{2}}{\sqrt{1-\cos \varphi}} \right] \frac{\pi}{2},$$

where $a = 10g \div 7(R+r)$.

This value of t , is not $[2 \div \sqrt{(2a)}] \log (1 - \sqrt{2})$, as indicated at p. 122, $[2 \div \sqrt{(2a)}] \log (1 - \sqrt{2})$ being the value of the integral at the superior limit only. At the inferior limit we have

$$\frac{\sqrt{1+\cos \varphi} - \sqrt{2}}{\sqrt{1-\cos \varphi}} = \frac{0}{0}, \text{ a vanishing fraction, whence}$$

$$\frac{-\sqrt{1-\cos \varphi}}{\sqrt{1+\cos \varphi}} = \frac{-0}{\sqrt{2}} = -0.$$

$$\therefore t = \frac{2}{\sqrt{(2a)}} \log \frac{1-\sqrt{2}}{-0} = \frac{2}{\sqrt{(2a)}} \log \frac{\sqrt{2}-1}{0} = \infty.$$

That is, the time required, to roll from the *summit* of the fixed sphere over one fourth of its circumference, is infinite.

If, instead of starting from the summit of the fixed sphere, the rolling sphere should start from a point, say $1''$ from the summit, the time, obviously, would be finite. In that case the time would be, as calculated by Mr. Kummell, $t = \sqrt{(2+a)} \times 12.7418368$.

SOLUTIONS OF PROBLEMS IN NUMBER FIVE.

SOLUTIONS of problems in No. 5 have been received as follows:

From R. J. Adcock, 277; Alex. S. Christie, 276, 278, 279, 280; Prof. W. P. Casey, 278; G. M. Day, 276, 278; Prof. E. J. Edmunds, 276, 277, 278; Henry Gunder, 276, 278, 280; Henry Heaton, 276, 277, 278, 279, 280; Chas. H. Kummell, 276, 277, 278, 279, 280; Prof. D. J. McAdam, 279; O. L. Mathiot, 276; W. L. Marcy, 276, 278, 279; K. S. Putnam, 276; P. Richardson, 276; Prof. E. B. Seitz, 276, 278, 279, 280; Prof. J. Scheffer, 278, 280; Prof. D. Trowbridge, 278, 279, 280.

276. "Given the chord AC of a circle, the side AB of a right angled triangle constructed on AC as hypotenuse, and the length of a perpendicular from A upon the line joining the right angle at B with the centre of the circle; to find the radius of the circle."